

RHINATHIOL Sanofi-Aventis

Composition

Carbocysteine (INN) 2 g/100 ml; syrup for children.
Also ADULT SYRUP 5 g/100 ml.

Also RHINATHIOL PROMETHAZINE

Carbocysteine 2g, promethazine hydrochloride
50 mg/100 ml syrup.

Indications

Acute and chronic bronchitis, rhinopharyngitis, tracheitis, acute or subacute laryngitis, otitis, sinusitis.

Contraindications

Adults: active peptic ulcer. Do not administer to infants of less than 1 year.

Side Effects/Precautions

Syrup for adult: When strong doses are given (4 x 15 ml spoonful) some digestive phenomena may occur: gastralgia, nausea, diarrhea.

Rhinathiol Promethazine syrup: drowsiness, the intake of alcohol during treatment is not recommended.

Dosage and Administration

Syrup for Children

Infants and children under 5 years: 1 x 5 ml spoonful once or twice a day.

Children over 5 years: 1 x 5 ml spoonful three times a day.

Syrup for Adults: 3 x 15 ml spoonful per day.

Promethazine Syrup

Adults: 3 tablespoons (3 x 15 ml)/day
Children over 5 years: 3 to 6 teaspoons (3 to 6 x 5 ml)/day.

Children under 5 years and infants over 1 year: 1 teaspoon (5 ml)/day/5 kg of bodyweight.

Packaging

sy: 125 ml (children)	125 ml
(adults)	125 ml
(Promethazine syrup).	